


RLINE Evaluation Database:

Raleigh 2006 NO Experiment



References

Baldauf, R., E. Thoma, A. Khlystov, V. Isakov, G. Bowker, T. Long, and R. Snow 2008: Traffic and Meteorological Impacts on Near-Road Air Quality: Summary of Methods and Trends from the Raleigh Near-Road Study. *Journal of the Air & Waste Management Association*, **58**, 865-878.

Venkatram, A., V. Isakov, E. Thoma, and R. Baldauf, 2007: Analysis of air quality data near roadways using a dispersion model. *Atmospheric Environment*, **41**, 9481-9497.

Files included

- Sources: "Ral2006_Source.txt"
- Receptors: "Ral2006_Receptors.txt"
- Meteorology: "Ral_2006_NO.sfc"
- RLINE input: "Line_Source_Inputs.txt"
- Data: "Raleigh_NOdata.xlsx" (contains measured traffic and concentrations)

Notes

- The line sources are run with unit emission rates so they can be multiplied by the traffic and emission factor to determine the modeled concentration.
- The line source is 1 km long, and 8 lanes are used (4 on each side of median).
- The emission factor used is 0.5 g/ veh./ km from Venkatram 2007.

- Each line of the met file represents a 10 minute period.

Run RLINE

Run RLINE using the provided “Line_Source_inputs.txt” file.

Interpreting Results

1. To determine the emission in g/m/s for each time period: multiply the traffic volume (in vehicle-miles) by the emission factor, convert mi. to km, divide by the length of the sources, divided by the number of seconds in your measurement time, and convert km to m.

$$(\text{unit emission rate}) \cdot \frac{\text{Traf. [veh. mi.]} \cdot 0.5 \left[\frac{g}{\text{veh. km}} \right] \cdot 1.6 \left[\frac{\text{km}}{\text{mi.}} \right]}{(8 \text{ lanes} \cdot 1 \text{ km}) \cdot 600 \text{ s} \cdot 1000 \left[\frac{\text{m}}{\text{km}} \right]} = \text{emission rate} \left[\frac{g}{\text{m} \cdot \text{s}} \right]$$

2. Multiply the unit-emission concentrations by the emission rate,

$$(\text{Output Conc.}) \cdot \left(\text{emission rate} \left[\frac{g}{\text{m} \cdot \text{s}} \right] \right) = \text{Conc.} \left[\frac{\mu g}{\text{m}^3} \right]$$

3. Divide by the density of NO and convert to ppb for comparison with the field data.

$$\frac{\text{Conc.} \left[\frac{\mu g}{\text{m}^3} \right]}{1.34 \left[\frac{\text{kg NO}}{\text{m}^3} \right]} \left[\frac{\text{kg}}{10^9 \mu g} \right] \cdot 10^9 = \text{Modeled Conc. [ppb NO]}$$