Application and verification of the NMMB/BSC-CTM forecast for solar energy





Albert Soret (1), Kim Serradell (1), Matthias Piot (2), Daniel Ortega (1), Vincenzo Obiso (1), Sara Basart (1) and Oriol Jorba (1) (1) Barcelona Supercomputing Center, Barcelona, Spain (albert.soret@bsc.es) (2) EnBW Energie Baden-Württemberg AG, Karlsruhe, Germany



In the beginning of April 2014, Northern Europe was affected by a mineral dust intrusion. On 4 April 2014, the power prediction for German solar installations was estimated as 21GW, whereas the measured power production merely reached 11GW. This strong overestimation significantly affected the hourly price in the wholesale electricity market: prices were firstly assessed at around 27€/MWh but rapidly reached a level close to 150€/MWh after recognizing the lack of solar output.

It has been found that a large proportion of the uncertainty of existing NWP models can be attributed to the lack of accurate aerosol data used in order to model solar radiation. Despite the advancements in the modelling of aerosol-cloud interactions, current meteorological models use parametrizations made mostly for climate considerations (generally monthly-based). The NMMB/BSC Chemical Transport Model (NMMB/BSC-CTM) is a new on-line chemical weather prediction system coupling atmospheric and chemical processes. The NMMB/BSC-CTM is the operational model of the Barcelona Dust Forecast Center of the WMO (http://dust.aemet.es). Previous studies have analyzed the predictability of the NMMB/BSC-CTM to reproduce dust outbreaks and their impact in solar radiation close to source regions such as the Mediterranean area and North Africa (Gkikas et al., 2015) where mineral dust has a high impact in the solar irradiance due to direct radiative effect.

The aim of this work is to analyze the performance of the NMMB/BSC-CTM to simulate a mineral dust intrusion far from the source regions to explore the applicability for solar energy forecast purposes.




Fig. 1. Backward trajectories ending at central Europe at 0.5, 5 and 10 km computed with HYSPLIT (Stein et al., 2015; Rolph, 20016)





Fig. 2. Aerosol Optical Depth at 550 nm and 10m wind speed: 28 March 2014 – 6 April 2014. NMMB/BSC-CTM 0.33° (RADON)

- Simulated period (spin-up period and key event): 20 March 2014 6 April 2014
- Key event: 28 March 2014 6 April 2014

NMMB/BSCDust model contains a dust module embedded online within the NCEP Nonhydrostatic Multiscale Model (NMMB). It provides weather and dust forecasts, from regional to global scales, thanks to its unified nonhydrostatic dynamical core. Dust cycle is represented through several parameterizations describing dust particles' sources, emissions, transport, removal from the atmosphere (wet and dry deposition) as well as the interaction with the radiation (Pérez et al., 2011).

| Table 1. Model description and main parameterizations | | | | | |
|---|--------------------|---------------------|-------------------------|----------------------------|------------------|
| Model | Spatial resolution | Vertical resolution | Met. initial conditions | Dust-radiation interaction | Land use |
| NMMB/BSC-CTM 0.1° (off) | 0.1° x 0.1° | 40 up to 50hPa | GFS | RADOFF | USGS/STATSGO-FAO |
| NMMB/BSC-CTM 0.33° (off) | 0.33° x 0.33° | 24 up to 50hPa | GFS | RADOFF | USGS/STATSGO-FAO |
| NMMB/BSC-CTM 0.33° (on) | 0.33° x 0.33° | 24 up to 50hPa | GFS | RADON | USGS/STATSGO-FAO |
| WRF-ARW v.3.7 | 12x12 km | 37 up to 50hPa | GFS | - | CORINE |



The NMMB/BSC-CTM model is able to reproduce the dust event. A comprehensive evaluation of this episode and other periods of study can be found here: http://www.bsc.es/earth-sciences/mineral-dust/nmmbbsc-dust-forecast/

Including a dust module in a model's calculations:

- improves its skill for reproducing surface irradiance during heavy dust events.
- under strong dust plume maximum temperature reduction (not shown).
- a higher resolution provides a higher mineral dust concentration, especially of dust peaks.
- differences between online/offline dust integration inside the model are moderate due to the relatively low dust concentrations in central Europe






Fig. 5. MODIS-Terra. Corrected reflectance, true color. 2014-04-04-10:40h




Fig. 6. Cloud cover. 2014-04-04-12:00h




Fig. 7. SW radiation at the ground. 2014-04-04-12:00h




Fig. 8. AOD at 550nm. 2014-04-04-12:00h

Ice-nucleation, cloud formation.

On 4 April 4th 2014, clouds covered central Europe (Fig. 5) however NWP models didn't predict such amount of clouds (Fig. 6 implying a significant overestimation of incoming radiation (Fig. 7). NMMB/BSC-CTM model properly

reproduced the dust event (Fig. 8), however the model didn't find any significant impact upon the incoming SW (Fig. 9).

Fig. 10 shows that the clouds observed in Fig. 5 are thin high-level clouds. The presence of mineral dust at high altitude during this period can suggest the formation of clouds due to the presence of mineral dust (ice-nucleation implementation; ongoing work).




Fig. 9. SW radiation at the ground evaluated against BSRN measurements (lefthand axis) in The Netherlands (cab station). AOD (right-hand axis).




Fig. 10. EUMETSAT MSG RGB. Dust appears pink or magenta. Dry land looks from pale blue (daytime) to pale green (nighttime). Thick, high-level clouds have red-brown tones and thin high-level clouds appear very dark.

Conclusions

The highest priority for energy network operators is the balance between energy demand and supply. Before the introduction of renewable energies, demand was matched with base-load power plants (coal and nuclear) and generating plants that can be scheduled, usually hydroelectric and fossil-based peak load power plants. This landscape has been changing rapidly with the integration of renewable energies into the energy mix over recent years.

Solar power forecasting prevents energy loss and improves the management of solar plants. In general terms, current NWP models don't use online predicted mineral aerosol concentration for radiation calculations. Central Europe is periodically affected by mineral dust intrusions, mainly in spring. Under these conditions, it is crucial to accurately model aerosol-cloud interactions.

The NMMB/BSC-CTM has proved to be a useful tool to predict mineral dust outbreaks:

- significant reduction of incoming SW under strong dust plumes.
- minor reductions for moderate events like the ones affecting central Europe.

On-going work

- Perform a climatological assessment to identify a statistically representative group of mineral dust episodes affecting central Europe to: 1) Analyze their impact on solar power generation; 2) Identify major weather events; 3) Analyze aerosol-cloud formation.; 4) Deposition
- Implementation and evaluation of indirect radiative effects.
- Integration of anthropogenic emission sources.

References and acknowledgment

- Gkikas et al. (2015): Atmospheric circulation evolution related to desertdust episodes over the Mediterranean. Q.J.R. Meteorol. Soc. Pérez et al. (2011): Atmospheric dust modeling from meso to global scales with the online NMMB/BSCDust model – Part 1: Model description, annual simulations and evaluation, Atmos. Chem. Phys., 11, 1300113027
- Stein et al., (2015): NOAA's HYSPLIT atmospheric transport and dispersion modeling system, Bull. Amer. Meteor. Soc., 96, 2059-2077.

Rolph, G.D. (2016): Real-time Environmental Applications and Display sYstem (READY) Website (http://www.ready.noaa.gov). NOAA Air Resources Laboratory, College Park, MD.

Acknowledgment: The authors gratefully acknowledge the NOAA Air Resources Laboratory (ARL) for the provision of the HYSPLIT transport and dispersion model used in this publication