



Analysis of the predictions of the Arctic sea-ice extremes in several forecasting systems

Neven S. Fučkar¹, Virginie Guemas^{1,2}, Matthieu Chevallier², Michael Sigmond³, Felix Bunzel⁴, and Francisco J. Doblas-Reyes^{1,5}

¹Institut Català de Ciències del Clima (IC3), Barcelona, Spain, ²Centre National de Recherches Météorologiques/Groupe d'Etude de l'Atmosphère Météorologique, Météo-France, CNRS, Toulouse, France, ³Candain Center for Climate Modeling and Analysis, Victoria, British Columbia, Canada, ⁴Max Planck Institute for Meteorology, Hamburg, Germany, ⁵Institució Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, Spain.



Sea Ice Concentration Anomalies
Sep 2014



● **Introduction and forecasting systems**

Since 1970s Arctic sea ice cover is undergoing substantial reduction and thinning in every month (especially September)

→ multi-year sea ice is being replaced with first-year sea ice making summer sea ice cover more variable from year to year and near-term climate prediction more challenging




We use a set of coupled climate forecast systems with full-field initialization:

- **EC-Earth2.3**
- **CNRM-CM (a successor of CM5)**
- **MPI-ESM**
- **CanCM3 and CanCM4**
Soon to include GloSea5 and CM2.1-FLOR seasonal forecasts


Forecast systems typically underestimate sea ice extent climatology

⇒ we need to remove mean bias, but also address conditional drift dependence

NH sea ice extent normal clim Nov 1 IC



NH sea ice extent normal clim May 1 IC



→ Forecast time (month)

- **Hierarchy of bias correction methods** (Fučkar et al., GRL 2014)

Even with the best possible forcing, BC and IC, model bias can be $>$ climate signal of interest

- 1) **Mean (per-pair) bias correction method**

→ replaces the long-term mean (over start dates i) of a model variable with the long-term mean of corresponding obs at each forecast time f (or lead time l)

Some forecast systems, beside the mean bias, also exhibit a conditional bias in time, i.e., a forecast drift that is dependent on the start date or other physically more relevant information



- 2) **Trend bias correction method**

→ replaces a linear regression of the forecasts on i with the linear regression of the corresponding obs on i at each forecast or lead time

Trend bias correction can also account for difference between the forecasted and obs. long-term trends, but even if they are similar there can be drift dependence on IC due to internal variability






- 3) **New IC bias correction method**

→ replaces a linear regression of monthly forecasts on OBS IC smoothed in time, e.g., obs in the first forecast month $o_{i,1}$, with the linear regression of monthly obs on $o_{i,1}$ for each forecast month f



Instantaneous IC at the beginning of a month are too noisy for the application in climate forecast on monthly and longer time scales
 ⇒ we need to use a smoothed proxy of obs IC (multiple options)






IC bias correction method has a strong sensitivity to the quality of obs ⇒ extending hindcast archive deeper into the past may not benefit the prediction skill (sparse and/or low quality obs), but with a shorter verification period we could increase the uncertainty



More sophisticated bias correction methods offer relevant but not statistically significant (95%) improvements over the mean (per-pair) adjustment method

95% confidence interval


→ Forecast time (month)



Trend bias correction method in analyzed models offers the most significant improvements in skill of the NH sea ice extent (SIE) in September likely due to significant errors in their long-term trends



→ **Forecast time (month)**

Trend bias correction method leads to an almost convergence of ACC in all models except one (initial shock issue?) \Rightarrow March and September are rather similar from point of anomaly correlation, but model errors ...




→ **Forecast time (month)**

Synoptic condition biases in summer and missing sea ice processes that can provide positive feedbacks (e.g., melt ponds, snow cover dynamics, etc.) are likely causing higher RMSE during boreal melting season than during boreal growing season




→ **Forecast time (month)**



NH sea ice extent

March


Nov 1 IC



NH sea ice extent

September


May 1 IC



NH sea ice extent


September

May 1 IC




↑ deviation of Arctic sea ice from obs trend can ↑ model errors in **summer**

Ensemble mean errors in **March** from Nov 1st IC



Ensemble mean errors in **September** from May 1st IC





→ Absolute distance of NH SIE obs from obs trend (M km²)

- **Summary and *future directions***

Choice of bias correction method has a substantial impact on deterministic prediction skill of the NH sea ice extent



ACC in March and September are similar, while September RMSE is significantly > than March RMSE



Winter extremes of NH SIE are better predicted than summer extremes that show biggest errors when the Arctic system is far from long-term trend in both positive and negative direction



- *Sep 2012 (,2007 & 2010?) case study*
- *Analysis of regional skill and its relation to structure of biases*

Anomaly correlation coefficients after applying **mean (per-pair) bias correction method**



→ **Forecast time (month)**

Root mean square errors after applying **mean (per-pair) bias correction method**




→ **Forecast time (month)**

NH sea ice extent

March


Nov 1 IC




NH sea ice extent

March

Nov 1 IC






NH sea ice extent


September

May 1 IC




Model errors ↑ in **summer** with ↑ deviation of Arctic sea ice from obs trend

Ensemble mean errors in **March** from Nov 1st IC





Ensemble mean errors in **September** from May 1st IC





→ **Absolute distance of NH SIE obs from obs trend (M km²)**

Anomaly correlation coefficients after applying **IC bias correction method**



→ **Forecast time (month)**

Root mean square errors after applying **IC bias correction method**



→ **Forecast time (month)**