

# MODELLING MINERAL DUST EMISSION OVER NORTH AFRICA AND MIDDLE EAST USING AN HIGH-RESOLUTION SURFACE PROPERTIES FROM SATELLITE RETRIEVALS



Sara Basart¹(sara.basart@bsc.es), Oriol Jorba¹, Carlos Pérez García-Pando<sup>2,3</sup>, Catherine Prigent⁴ and José María Baldasano<sup>1,5</sup>

1 Earth Sciences Department, Barcelona Supercompuling Center-Centro Nacional de Supercomputación (BSC-CNS), Barcelona, Spain litute at Columbia University, NASA Goddard Institute for Space Studies and International Research Institute for Climate and Society, New York, USA
2 Department of Applied Physics and Applied Math, Columbia University, New York, New York, USA
4 Laboratolier de Etudes du Rayonnement et de la Matière en Astrophysique, CNRS UMR 8112, Observatoire de Paris, Paris, France
1 Environmental Modelling Laboratory, Technical University of Catalonia (UPC), Barcelona, Spain

### **OBJECTIVE**

Implement and test the impact of the drag partition correction in the emission scheme of the NMMB/BSC-Dust model using the aerodynamic roughness length from ASCAT/PARASOL

### INTRODUCTION

The threshol friction velocity  $(u_{*th})$  is defined as the velocity above which soil particles begin to

Threshold friction velocity over smooth surface and  $D_p$  the diameter of soil particles



for the supression of soil erosion in wet soils

**Drag partition correction** Z<sub>o</sub>: roughness length

Z<sub>0s</sub>: roughness length over smooth surfaces

The *drag partition correction (R)* accounts for the presence of nonerodible roughness elements (rocks, pebble, vegetation, etc.) in natural land surfaces. Aeolian aerodynamic roughness length (z<sub>0</sub>) in arid regions is a key parameter to predict the vulnerability of the surface to wind erosion and the related production of mineral aerosol.

### **TOOLS**

### NMMB/BSC-Dust

The NMMB/BSC-Dust model is a part of the NMMB/BSC-Chemical Transport Model (NMMB/BSC-CTM; Pérez et al. 2011; Jorba et al., 2012; Spada et al., 2013) a new chemical

- weather prediction system. Its main features are: It allows working in global and regional configuration
  - A physically emission scheme, which includes saltation and sandblasting
  - 8 bins externally mixed
  - Wet removal in-cloud and below-cloud scavenging

The NMMB/BSC-Dust provides operational daily dust forecasts at regional and global domains and participates in:



### ASCAT/PARASOL

Satellite-derived roughness length at the global scale have emerged and provide the opportunity to use them in advanced emission schemes. A global map of the aeolian aerodynamic roughness length at high resolution is derived, for arid and semi-arid regions merging PARASOL and ASCAT data (Prigent et al., 2012).





- Z<sub>0</sub> derived from ASCAT backscattering (1999-2009) and PARASOL 865 nm (2005-2008)
- Spatial resolution: ~ 6km x 6km , (1/100º x 1/18º) Valid range: 0 - 0.08 cm. They are
- calculated outside this range but their validity is questionable.

### **CONCLUSIONS**

This contribution shows the analysis of the impact of the drag partition correction in the emission scheme of the NMMB/BSC-Dust model using the aerodynamic roughness length from ASCAT/PARASOL. These previous results highlight

- Over Sahara, the results from ASCAT/PARASOL are comparable to LISA. Moreover, the high minimum value of the NMMB roughness length (at  $z_0$  = 0.1001m) I inhibits dust emissions. M04 provides higher emissions than MB95 as expected. M04 modified MB95 to accommodate the higher roughness
- length in the case of vegetated surfaces.
- DRAG (ASCAT/PARASOL-MO4) provides higher AOD values than NO-DRAG in winter and spring in the Sahara.

  Otherwise, in NW. Africa, AOD fields are lower with DRAG (ASCAT/PARASOL-MO4) than NO-DRAG. This is due to the impact of the NMMB roughness length in this region.
- In the AERONET annual comparison, NO-DRAG shows slightly better skill scores than DRAG. However, both model  $configurations\ present\ similar\ tendencies\ in\ the\ different\ regions\ analysed.$

Ongoing simulations at different spatial resolutions and with a refined roughness length map are under analysis.

### 1. SENSITIVITY EXPERIMENTS

- The simpler default option of the atmospheric model NCEP-NMMB applies a constant  $z_0$  over desert regions ( $z_0 = 0.1001m$ )
- To asses the validity of the ASCAT/PARASOL data, results is compared with those obtained with LISA dataset (Laurent et al., 2005) which provides an estimate of z<sub>0</sub> at 0.25ºx0.25º spatial resolution for arid and semi-arid areas of Africa and Asia.
- Additionally, two different approaches to account for the drag partition correction (R) been tested: Marticorena and Bergametti (1995; referred as MB95) and MacKinnon et al. (2004; referred as M04).
- They are compared with the operational model version that no includes the drag partition correction and includes a topographical preferential source mask from Ginoux et al. (2001) (referred as NO-DRAG)










### RESULTS: 26th March to 19th April 2011

nulated emissions (g/m²) from 26<sup>th</sup> March to 19<sup>th</sup> April




### AERONET Level 2.0 comparison



### 2. ANNUAL MODEL EVALUTION

### Seasonal averages for 2011



Both model configurations, DRAG (ASCAT/PARASOL-M04) and NO-DRAG, reproduces the seasonal evolution observed also in MISR.

Temporal period: 2011
Regional domain: 31°W-71°E and 0°-70°N
Spatial resolution: 0.25° x 0.25° x 40 layers

conditions from NCEP/FNL rological analysis (19x19) at OUTC ar ary conditions every 6 h444

- In summer, low AOD values in both version: are linked to the initial meteorological fields
- are linked to the initial meteorological fields from NCEP/FNL.

  DRAG provides higher AOD values over the Sahara-Sahel region as well as Middle East. On the contrary, in NW. Africa NO-DRAG shows higher AOD values because the impact of the NMMB roughness length.

## **AERONET comparison for 2011**



- 49 stations distributed in 9 region

•	AE < 0.75 is considered in the calculation
•	AE >= 1.2 we assign AOD observe = 0

Regions	NDATA	REG NO-DRAG			REG DRAG ASCAT/PARASOL MO4		
		r	RMSE	MB	r	RMSE	MB
Sahara-Sahel	3337	0,56	0,36	-0,05	0,49	0,52	0,04
E.Trop.N.Atl	2218	0,70	0,26	-0,17	0,58	0,32	-0,14
E.sub-Trop.N.Atl	1274	0,72	0,19	-0.12	0,62	0,22	-0,14
NW. Africa	2254	0,76	0,10	-0,05	0,70	0,12	-0,06
WJP	4941	0,76	0,09	-0,03	0,73	0,10	-0,04
E.IP-W.Med	4971	0,69	0,07	-0,01	0,63	0,07	-0,02
C. Med	5595	0,67	0,08	-0,01	0,54	0,10	-0,02
E. Med	4176	0,54	0,12	-0,01	0,53	0,12	-0,03
West Asia	1240	0,37	0,43	-0,30	0,29	0,49	-0,34
NAMEE	29446	0,75	0,19	-0,05	0,66	0,24	-0,05

### References

Pérez et al., 2011: Almospheric duat modelling from meso to global scales with the online NMMB/BSC-Duat model - Part 1: Model description, annual simulations and evaluation. Almos. Chem. Phys., 11.
Jorba et al., 2012: Pédratial significance of photeexcled NO2 on global air quality with the NMMB/BSC chemical transport model, J. Geophys. Res., dots 10/29/2012/201730.

Spadul et al., 2013: Modelling and evaluation of the global see-seat aerosol distribution: sensitivity to size-resolved and sea-surface temperature dependent emission schemes, Almos. Chem. Phys., 13, 11735-11755.

Matricorna and Bergamett, 1995: Modeling the attriospheric dust cycle: 1. Design of a soli-derived dust emission scheme, J. Geophys. Res., 100, 4415-14420.

4145-1430.
MacKimon et al. 2004: Comparison of aerodynamically and model-derived roughness lengths (zo) over diverse surfaces, central Migiave Desert,
Zalfornia, USA: Geomorphology 63 (2004) 103-113.
"Yighert et al. 2012 Comparison of saltelle microwave backscattering (ASCAT) and visible/ineas-infrared reflectances (PARASOL) for the estimation of

### Acknowlegments

The authors acknowledge the CICYT project (CGL2013-46736) and Severo Ochoa (SEV-2011-00067) programme of the Spanish Government. Simulations have been performed in the Murenostrum supercomputer. The authors thank AERONET, MISR, principal investigators and scientists for establishing and maintaining the AERONET and MISR data.