







INTAROS plans for model evaluation

Ralf Doescher (SMHI), Yongqi Gao (NERSC), *Virginie Guemas* (*BSC*), Stein Sandven (NERSC)









Improving skill of climate predictions

Goal: better prediction skill

Input: **observations**, via reanalysis, either by "big" actors (ECMWF) or by individual institutes, with or without INTAROS data.

Means: improved accuracy and suitability of **initial conditions** in combination with global climate models and Arctic hydrological models.

T/S profiles or gridded (at least SST/SSS), sea ice concentrations.

Expected results: more skilful climate prediction.

Formal distribution in 3 activities

- Understand dependence of climate simulations and variability on data availability
- 2) Assessing the full effect of enhanced initialization on climate prediction skill, and promote INTAROS data in the climate prediction community
- 3) Improving hydrological runoff predictions






Member1

Member2

Member29

Norwegian Climate Prediction Model (NorCPM)





V0 system :**SST**

V1 system: SST+T-S (+SSH?)


V2 system: **Ice-concentration**

V3 system: All



Propagation

correction










Ec-Earth












Experimental setup



1993










Experimental setup












Experimental setup
























INTAROS model evaluation workplan

- 1. Arctic climate prediction with NorCPM and EC-Earth 3.2 by assimilating the additional data (sea ice, sea surface temperature) from INTAROS. The aim is to test the added-value of INTAROS data set.
- 2. Data-deny experiments (under planning)
- 3. Improve hydrological runnoff predictions.