



**Barcelona
Supercomputing
Center**
Centro Nacional de Supercomputación



A source apportionment assessment of O₃ in peak summer events over southwestern Europe


María Teresa Pay , Carlos Pérez-García Pando,
Marc Guevara, Sergey Napelenok, Xavier
Querol

28/11/2017

26th GLOREAM Meeting - Berlin (Germany)

Background and Motivation


O₃ dynamic



Sources:

Millán et al., 1997;2000, 2014; Gangoiti et al, 2001, 2002, 2006
Toll and Baldasano, 2000

O₃ Trends and exceedances




Source: Querol et al. (2016 AE).

Open questions


- What are the sources responsible for the high O₃ concentration over the whole Spanish Peninsular territory?
- Can administrations implement control strategies that are effective to reduce high O₃ concentration?

Source Apportionment (SA) methods in AQM

Zero-Out (ZO)



SA within the AQM



Advantages

- Time saving (one simulation)
- Mass consistency
- Real atmospheric conditions
- Fully traceable

Objective

Unraveling the origin of the high surface O₃ concentrations in the Spanish Iberian Peninsula (IP)

- Quantifying the contribution from:
 - the **main NO_x emission sectors within the IP**
 - the **external contribution** (O₃ produced outside of the IP)
- Using the Integrated Source Apportionment Method (ISAM) in the CALIOPE air quality modelling System at high resolution over the IP.



The CALIOPE System



Set-up for the Source Apportionment

METEO

- **WRF-ARWv3.5** (RRTM/WSM3/YSU/NoahLSM)
- Ver. Res.: 37σ / 50hPa (top)
- Hor. Res: **12 km (EU12) 4 km (IP4)**
- **IC/BC (EU12/IP4): GFS (NCEP) / nesting EU12**

EMIS

- **HERMESv2.0**
- EU12: HERMES-DIS (EMEP data 2009)
- **IP4: HERMES-BOUP (Spain) + HERMES-DIS(Europe)**
- Biogenic emission MEGANv2.0.4

CHEM

- **CMAQv5.0.2** (**ISAM**, CB05TUCL, AERO6)
- Ver. Res: 37σ / 50hPa (top)
- Hor. Res: 12 km (EU) 4 km (IP)
- BC (EU12/IP4): MOZART4-GEOS-5/nesting EU12
- MCIPv4.0



O₃ Integrated Source Apportionment Method (ISAM)

- Augmented version of **CMAQv5.0.2** (AERO6, CB05TUCL) (*Kwok et al., 2013; 2014*)
- O₃ formation regime (NO_x- or VOC-limited conditions): **ratio H₂O₂/HNO₃** (*Zhang et al., 2009*).

$$P_{tag}^{N,new} = P_{tag}^{N,old} + P_{bulk}^{new} \frac{\sum_x NO_{x,tag}}{\sum_{tag} \sum_x NO_{x,tag}}$$

NO_x-limited conditions - Ratio_{H₂O₂/HNO₃} > 0.35

$$P_{tag}^{V,new} = P_{tag}^{V,old} + P_{bulk}^{new} \frac{\sum_y VOC_{y,tag} \times MIR_y}{\sum_{tag} \sum_y VOC_{y,tag} \times MIR_y}$$

VOC-limited conditions - Ratio_{H₂O₂/HNO₃} < 0.35

NO_{x,tag}: concentrations of the nitrogen species in CB05 (9 species)

VOC_{j,tag}: concentrations of the VOC species in CB05 (14 species)


MIR_y: Maximum Incremental reactivity factor of the VOC species y, corresponding to the O₃ generating potential of each single VOC specie.

- The bulk O₃ concentration in each model grid cell (P_{bulk}) is equal to the sum of O₃ tracers that were produced in either NO_x or VOC-sensitive conditions

$$P_{bulk} = \sum_{tag} P_{tag} = \sum_{tag} P_{tag}^N + \sum_{tag} P_{tag}^V \quad \text{Mass conservative}$$

O₃ tagged sources in this study

Annual emissions HERMESv2.0 in Spain 2009






ISAM tag*	Emission by SNAP category	Description
SNAP1	SNAP1	SNAP1: Energy industry
SNAP34	SNAP34	SNAP34: Industry (combustion and processes)
SNAP7	SNAP7	SNAP7: Road transport, exhaust and non-exhaust
SNAP8	SNAP8	SNAP8: Non-road transport (international shipping)
OTHR	SNAP2 + SNAP5 + SNAP6 + SNAP9 + SNAP10 + SNAP11	SNAP2: residential and commercial/institutional combustion SNAP5: Fugitive emissions from fuels SNAP6: Product use including solvents SNAP9: waste management SNAP10: Agriculture SNAP11: Other sinks
BCON	-	Chemical boundary conditions to IP4 domain from the EU12 simulation which includes the contribution from Europe and international contribution from MOZART-4
ICON	-	Initial chemical condition of the domain IP4



• <1 %

O₃ episode: July 21-31 2012







Meteorological context (6 UTC)

July-25





July-28

July-31





W10 / T2M



Prec. / MSLP





Geo. Height / T




ITL and NWadv
represent 44% of the
days in the IP both
taking place in
summer
(Valverde et al., 2015)

Source-sector contribution during peaks: ITL – July 25

BSC-ES/AQF ARWv3+CMAQ-ISAM+HERMESv2 O3 ($\mu\text{g}/\text{m}^3$)
p90 forecast for 25 Jul 2012 - Res:4x4km




BSC-ES/AQF ARWv3+CMAQ-ISAM+HERMESv2 O3 SNAP1 ($\mu\text{g}/\text{m}^3$)
p90 forecast for 25 Jul 2012 - Res:4x4km




BSC-ES/AQF ARWv3+CMAQ-ISAM+HERMESv2 O3 SNAP34 ($\mu\text{g}/\text{m}^3$)
p90 forecast for 25 Jul 2012 - Res:4x4km




BSC-ES/AQF ARWv3+CMAQ-ISAM+HERMESv2 O3 SNAPB ($\mu\text{g}/\text{m}^3$)
p90 forecast for 25 Jul 2012 - Res:4x4km




BSC-ES/AQF ARWv3+CMAQ-ISAM+HERMESv2 O3 SNAP7 ($\mu\text{g}/\text{m}^3$)
p90 forecast for 25 Jul 2012 - Res:4x4km



BSC-ES/AQF ARWv3+CMAQ-ISAM+HERMESv2 O3 OTHR ($\mu\text{g}/\text{m}^3$)
p90 forecast for 25 Jul 2012 - Res:4x4km




BSC-ES/AQF ARWv3+CMAQ-ISAM+HERMESv2 O3 BCON ($\mu\text{g}/\text{m}^3$)
p90 forecast for 25 Jul 2012 - Res:4x4km




10.5°W 9°W 7.5°W 6°W 4.5°W 3°W 1.5°W 0° 1.5°E 3°E 4.5°E

10.5°W 9°W 7.5°W 6°W 4.5°W 3°W 1.5°W 0° 1.5°E 3°E 4.5°E




Regionalization of source-sector contribution



- CIP
- EIP
- EV
- GV
- MED
- NEIP
- NIP
- NWIP
- SIP
- WIP


Daily mean contribution during
DMA8 > 120 μm^3



Evaluation

EIONET Spanish monitoring stations: 347 O₃ and 357 NO₂ (85% temporal coverage in the episode)


- O₃ at RB (±4 µg/m³) and r > 0.6 (50% stations).
- NO₂ highest underestimation at TR (-7 µg/m³) and r >0.6 (25 % stations.)




MB (µg/m ³)		O ₃ DMA8				
		<-40	(-40,-10]	(-10,10]	(10,40]	>40
O ₃ HL	<-40	0	0	0	0	0
	(-40,-10]	0	7 (2%)	0	0	0
	(-10,10]	0	A 35 (10%)	B 94 (28%)	5 (2%)	0
	(10,40]	0	3 (1%)	C 65 (19%)	D 122 (36%)	5 (2%)
	>40	0	0	0	1 (<1%)	5 (2%)

- | | | | |
|-------------|--------------|------------|--------------|
| 1 - CIP - D | 3 - NEIP - B | 5 - GV - D | 7 - NWIP - D |
| 2 - CIP - B | 4 - NEIP - B | 6 - GV - D | 8 - NWIP - B |




Central Iberian Peninsula (CIP)



(a) 1 - CIP - D - ES0126A - BU




(b) 2 - CIP - B - ES1537A - BU




- sa.fac
- ICON
 - BCON
 - OTHR
 - SNAP1
 - SNAP34
 - SNAP7
 - SNAP8
- ref.fac
- obs
 - cmaq


Northeastern Iberian Peninsula (NEIP)



(c) 3 - NEIP - B - ES1992A - BU




(d) 4 - NEIP - B - ES1778A - BR




- sa.fac
- ICON
 - BCON
 - OTHR
 - SNAP1
 - SNAP34
 - SNAP7
 - SNAP8
- ref.fac
- obs
 - cmaq


Gualdaquivir Valley (GV)



(a) 5 - GV - D - ES1644A - BU




(b) 6 - GV - D - ES1793A - BR




- sa.fac
- ICON
 - BCON
 - OTHR
 - SNAP1
 - SNAP34
 - SNAP7
 - SNAP8
- ref.fac
- obs
 - cmaq


North Western Iberian Peninsula (NWIP)



(c) 7 - NWIP - C - ES1868A - BU



(d) 8 - NWIP - B - ES0005R - BR



- sa.fac
- ICON
 - BCON
 - OTHR
 - SNAP1
 - SNAP34
 - SNAP7
 - SNAP8
- ref.fac
- obs
 - cmaq

Conclusions

- The **external contribution** accounted $> 45\%$ of the O_3 under exceedances of the $120 \mu\text{g m}^{-3}$ threshold for the DMA8 O_3 (all regions).
 - Downward mixing of O_3 upper layers transported from beyond the IP in CIP.
 - Recirculation in the Spanish Mediterranean coasts.
- **Contribution from local/regional sources** is significant in O_3 peaks downwind of NO_x hotspots.
 - Central and NE IP (big cities in Spain): the highest **road transport** contribution to O_3 (up to 40% in daily peak during events).
 - Industrial regions (N and NW IP and Guadalquivir Valley): **energy generation and industrial processes** contribute to O_3 up to 11%.
- The **non-road transport** is a contributor as significant as the road transport in all sub-regions (10-19%).
- **ISAM-CALIOPE** useful tool:
 - Source contribution assessment in the Spanish IP.
 - Identification of potential errors in emission estimates from different sectors: shipping, traffic emissions.
 - Necessities of improvement in meteorology under-stagnant conditions.
 - Design more cost-efficient mitigation plans (together with source sensitivity).



**Barcelona
Supercomputing
Center**
Centro Nacional de Supercomputación




EXCELENCIA
SEVERO
OCHOA

THANK YOU!


maria.pay@bsc.es

www.bsc.es


O3 cross sections




BSC-ES/AQF ARWv3+CMAQ-ISAM+HERMESv2 O3 BCON ($\mu\text{g}/\text{m}^3$)
p90 forecast for 25 Jul 2012 - Res:4x4km



BSC-ES/AQF ARWv3+CMAQ-ISAM+HERMESv2 O3 BCON ($\mu\text{g}/\text{m}^3$)
p90 forecast for 28 Jul 2012 - Res:4x4km




BSC-ES/AQF ARWv3+CMAQ-ISAM+HERMESv2 O3 BCON ($\mu\text{g}/\text{m}^3$)
p90 forecast for 31 Jul 2012 - Res:4x4km




Source-sector contribution during peaks: Nwad – July 28


BSC-ES/AQF ARWv3+CMAQ-ISAM+HERMESv2 O3 ($\mu\text{g}/\text{m}^3$)
p90 forecast for 28 Jul 2012 - Res:4x4km




BSC-ES/AQF ARWv3+CMAQ-ISAM+HERMESv2 O3 SNAP1 ($\mu\text{g}/\text{m}^3$)
p90 forecast for 28 Jul 2012 - Res:4x4km




BSC-ES/AQF ARWv3+CMAQ-ISAM+HERMESv2 O3 SNAP34 ($\mu\text{g}/\text{m}^3$)
p90 forecast for 28 Jul 2012 - Res:4x4km




BSC-ES/AQF ARWv3+CMAQ-ISAM+HERMESv2 O3 SNAPB ($\mu\text{g}/\text{m}^3$)
p90 forecast for 28 Jul 2012 - Res:4x4km






BSC-ES/AQF ARWv3+CMAQ-ISAM+HERMESv2 O3 SNAP7 ($\mu\text{g}/\text{m}^3$)
p90 forecast for 28 Jul 2012 - Res:4x4km



BSC-ES/AQF ARWv3+CMAQ-ISAM+HERMESv2 O3 OTHR ($\mu\text{g}/\text{m}^3$)
p90 forecast for 28 Jul 2012 - Res:4x4km



BSC-ES/AQF ARWv3+CMAQ-ISAM+HERMESv2 O3 BCON ($\mu\text{g}/\text{m}^3$)
p90 forecast for 28 Jul 2012 - Res:4x4km




OTHER

BCON


Source-sector contribution during peaks

ITL – July


BSC-ES/AQF ARWv3+CMAQ-ISAM+HERMESv2 O3 ($\mu\text{g}/\text{m}^3$)
p90 forecast for 31 Jul 2012 - Res:4x4km




BSC-ES/AQF ARWv3+CMAQ-ISAM+HERMESv2 O3 SNAP1 ($\mu\text{g}/\text{m}^3$)
p90 forecast for 31 Jul 2012 - Res:4x4km




BSC-ES/AQF ARWv3+CMAQ-ISAM+HERMESv2 O3 SNAP34 ($\mu\text{g}/\text{m}^3$)
p90 forecast for 31 Jul 2012 - Res:4x4km



BSC-ES/AQF ARWv3+CMAQ-ISAM+HERMESv2 O3 SNAP8 ($\mu\text{g}/\text{m}^3$)
p90 forecast for 31 Jul 2012 - Res:4x4km






BSC-ES/AQF ARWv3+CMAQ-ISAM+HERMESv2 O3 SNAP7 ($\mu\text{g}/\text{m}^3$)
p90 forecast for 31 Jul 2012 - Res:4x4km




BSC-ES/AQF ARWv3+CMAQ-ISAM+HERMESv2 O3 OTHR ($\mu\text{g}/\text{m}^3$)
p90 forecast for 31 Jul 2012 - Res:4x4km




BSC-ES/AQF ARWv3+CMAQ-ISAM+HERMESv2 O3 BCON ($\mu\text{g}/\text{m}^3$)
p90 forecast for 31 Jul 2012 - Res:4x4km









BSC-ES/AQF ARWv3+CMAQ-ISAM+HERMESv2 O3 SNAP7 ($\mu\text{g}/\text{m}^3$)
00h forecast for 00UTC 31 Jul 2012 - Res:4x4km



BSC-ES/AQF ARWv3+CMAQ-ISAM+HERMESv2 O3 BCON ($\mu\text{g}/\text{m}^3$)
00h forecast for 00UTC 31 Jul 2012 - Res:4x4km





STATE vs RESPONSE

What are the various contributors to modeled concentrations?

Source contribution approaches: tracks the formation and transport of O_3 and $PM_{2.5}$ from specific sources and allows the calculation of contributions at each hour and grid cell

STATE: relative importance of sources that contribute to high concentrations)

- ✓ Brute force zero out
- ✓ Ozone and PM source apportionment



How will the modeled concentrations change based on changes to emissions?

Source sensitivity approaches: estimates sensitivity coefficients that relate emission changes from specific emission sources to model outcomes at each hour and grid cell

RESPONSE: prediction of how pollutant will respond to reductions in precursor emissions

- ✓ Brute force zero out
- ✓ Decoupled Direct Method (DDM)

DDM compared to Brute Force

