

# Development of a wind energy climate service based on seasonal climate predictions



Verónica Torralba, Francisco J. Doblas-Reyes, Isadora Jiménez, Llorenç Lledó, Nube González-Reviriego and Albert Soret  
 Barcelona Supercomputing Center (BSC), Catalan Institute of Climate Science (IC3), Institució Catalana de Recerca i Estudis Avançats (ICREA)


## Climate information

The large amount of information that arises from the climate is hard to understand and in most cases wind energy users are not able to incorporate it in their daily activities. The main goal of this work is the creation of tailored climate information that can be afterwards used as a tool to inform wind energy users with greater accuracy than their current approaches.

1


## Wind speed assessment

Wind energy users can not use the outputs of the global climate models directly because they require an evaluation and interpretation of the forecast quality




Range of the differences (m/s) between the 10-m wind speed values produced by ERA-Interim, MERRA and JRA-55. in DJF.

Verification of seasonal forecasts against different reanalyses might have slightly different results related with the reanalyses uncertainty.



Correlation of 10-m wind speed between the ensemble mean forecasts from ECMWF S4 and ERA-Interim reanalysis in DJF

Key regions for the wind industry as North America, Northeast Brazil, North Sea and Eastern China display potential skill.



Bias for 10-m wind speed between the seasonal forecasts from ECMWF S4 and ERA-Interim reanalysis in winter (DJF)

Biases resulting from the prediction system inability to perfectly reproduce the climate variability should be corrected in order to produce usable climate information.

2

## Bias adjustments


Different methods of bias-correction have been used to produce forecasts with improved statistical properties

Method	Equation	Description	Result
Simple bias correction	$y_{j,i} = (x_{ij} - \bar{x}) \frac{\sigma_{ref}}{\sigma_e} + \bar{o}$	Based on the assumption that both the reference and forecasted distribution are well approximated by a Gaussian distribution.	
Calibration method	$y_{j,i} = \alpha x_i + \beta z_{ij}$	The variance inflation modifies the predictions to have the same interannual variance as the reference dataset and corrects the ensemble spread to obtain more reliable probabilities.	
Quantile mapping	$y_{j,i} = (ecdf^{ref})^{-1} ecdf^{mod}(x_{ij})$	Each forecast is assigned to the quantile of the forecast climatology it corresponds, and then the forecast value is changed by the observed value in the same quantile of the of the observational climatology.	


3

## Forecast quality assessment

The impact of the bias-adjustments on the forecast quality has been explored by several scoring measures



- Raw forecasts display poor reliability.
- Corrected forecasts display improved reliability with more points falling along the diagonal and more homogeneously populated sharpness diagrams.
- Calibrated forecasts show slopes closer to one.







Calibration method produces the best CRPSS and reliability for the below-normal category and above normal category, however the forecasts corrected with the quantile mapping method produce higher RPSS.

4

## Transfer model

Capacity factor (CF) is widely used indicator for the wind energy users. Predictions of CF are not produced by the climate models and it should be computed from climate variables



## Wind energy decisions

Climate predictions tailored to the wind energy sector represent an innovation to better understand the future variability of wind energy resources. These predictions can improve decision making processes related with: maintenance works, the matching of supply with demand and the reduction of financial penalties for incorrect wind power predictions.

Development of a wind energy climate service