



---

# AECT-2017-2-0009 Volcanic activity in seasonal to decadal climate forecasts

---

## 1. General Information

---

### Activity Id

AECT-2017-2-0009

### a) Activity Title

Volcanic activity in seasonal to decadal climate forecasts

### b) Area

Astronomy, Space and Earth Sciences

## 2. Research Project Description

---

### a) Is this a Test Activity?

No

### b) Is this a Long Term Activity that will extend over two application periods?

Yes

### c) Brief description of the Project

At the intermediate timescales between weather forecasts and climate predictions, seasonal to decadal forecasting represents a challenge that must be met both for increased scientific understanding and to develop societal applications, in particular for agriculture and energy production and consumption. The Barcelona SuperComputing Center (BSC) is the only Spanish institute that is currently able to provide such forecasts in a real-time context, using all available sources of climate predictability and shaping forecast systems for services. Over the last years, the Earth Department of the BSC illustrated the predictability associated with the initialisation of climate models with the best available observations, which could be seen both in terms of global temperature and regional climate variability. An open aspect of this activity is linked to the role of the stratospheric particles that strongly impact the climate variability from intra-annual to inter-annual timescales through a complex dynamical response of the atmosphere to aerosol forcing. Stratospheric particles mainly originate from

volcanic eruptions. Martin Ménégoz is coordinating at BSC the “Volcanic activity in seasonal to decadal climate forecasts” project, (VOLCADEC, Ministerio de Competitividad y Economía (MINECO), 168 K€) with the aim to answer how forecast systems should take into account these particles. It focuses on two main challenges: (1) how can the climate response to a large volcanic eruption and to stratospheric aerosols changes be predicted? (2) How climate conditions modulate the climate response to large volcanic eruptions? The final goal of VOLCADEC is to design the optimal strategy to take into account stratospheric aerosols in climate forecast systems, during and outside periods of volcanic eruptions. Optimized systems will be helpful to provide climate information, in particular in the Mediterranean region, where the climate variability is largely driven by stratospheric aerosols.

This request of computing time to RES involves extensive modelling using the EC-Earth climate model to contribute to the VOLCADEC project. At the international level, these experiments will be part of the “Model Intercomparison Project on the climatic response to Volcanic forcing to investigate the climate response to volcanic eruptions” (VOLMIP, <http://volmip.org/>) and the “Decadal Climate Prediction Project” (DCPP, <https://www.wcrp-climate.org/dcp-overview>), two exercises focusing on the ability of current climate models to produce seasonal to decadal forecasts. Contributing to these projects is an opportunity for the Earth department of the BSC to reinforce its involvement in CMIP6 (<https://www.wcrp-climate.org/wgcm-cmip/wgcm-cmip6>), the international scientific initiative that is planned beforehand the writing of the next IPCC report (<https://www.ipcc.ch/>).

#### **d) Grant References**

Project VOLCADEC MINECO/FEDER (ref. CGL2015-70177-R)  
European project SPECS FP7 308378

#### **e) Brief description of the Project (if this Activity takes place in the context of a Technology or Industrial Project)**

#### **f) Specific Activity proposed**

Introduction

Apart from human factors impacting the climate with a response time of several decades, natural variability induces a strong inter-annual to decadal climate variability that is challenging to forecast (Hawkins et al., 2009). There is a strong need for a better understanding of the natural variability to design models able to provide forecasts for the next years to decades. Climate models still need significant progress for successful operational applications at seasonal to decadal scales. Initialisation of climate models is fundamental to forecast temperature and precipitation at these time scale (Doblas-Reyes et al., 2013), but external radiative forcings play also a significant role: variations of greenhouse gases, human-made aerosols, volcanic particles, solar irradiance and land use changes are the main forcings which impact climate decadal variability (e.g. Hansen et al., 2011). Among these factors, volcanic aerosols - the main source of stratospheric aerosols - have inter-annual variations whose impact is challenging to estimate because their representation in climate models is generally basic. As a consequence, it is difficult to estimate the predictability associated to stratospheric aerosols in seasonal to decadal forecasts. In addition, when the next large volcanic eruption will occur, the uncertainties on aerosols

emissions added to the low skill of current climate models will strongly limit our possibility to forecast its seasonal to decadal climatic impacts.

Agung (1963), El Chichón (1982) and Pinatubo (1991) are the last three major eruptions with significant climatic impact. By injecting a large amount of sulphate aerosols into the stratosphere, such eruptions partly blocked the downward solar radiation, inducing a cooling of the troposphere ranging between 0.1°C and 0.5°C. However, it is challenging to understand the regional climate response to these eruptions that is often driven by complex dynamical processes involving both the ocean and the atmosphere (Swingedouw et al, 2017). Due to the low number of observations of both the eruptions and their climate impacts, it is difficult to isolate its signal from the high natural variability of the climate system, and in particular its interactions with the El Niño Southern Oscillation (ENSO) and the North Atlantic Oscillation (NAO), the latter strongly driving temperature and precipitation rates in the Mediterranean area (Bladé et al., 2011). Furthermore, recent investigations showed that small eruptions also affect the stratospheric aerosol burden and could have explained part of the current slowdown of the global warming (Solomon et al., 2011). In a general manner, there is a strong need to design climate models able to forecast the climate response to stratospheric aerosols changes at seasonal to decadal timescales. This is the aim of the VOLCADEC project for which we apply for computation hours to the Red Española de Supercomputación (RES).

People involved in the project

The VOLCADEC project is carried out within the Climate Prediction team (<https://www.bsc.es/discover-bsc/organisation/scientific-structure/climate-prediction>) of the Barcelona Supercomputing Center (BSC), involving Martin Ménégos (PI of the project, with 17 publications focusing on climate variability and aerosols), Roberto Bilbao (expert in oceanography), Omar Bellprat (expert in statistical analysis of climate data), Virginie Guemas (head of the climate prediction group) and Francisco Doblas-Reyes (head of the Earth Department at BSC). The VOLCADEC research is also supported by the overall climate prediction team that develops statistical tools and invests heavily in the development of the EC-Earth climate model (Hazeleger et al., 2010). The VOLCADEC team includes also two internationally recognized French researchers: Christophe Cassou (CNRS, France, expert in decadal climate variability) and Didier Swingedouw (CNRS, France, expert in climate response to volcanoes). The VOLCADEC activities are also related to several European projects, in particular the Seasonal to Decadal climate Predictions for the improvement of the European Climate Services (<http://www.specs-fp7.eu/>), recently led by BSC and bringing together the 20 main European institutes investigating the climate predictability at seasonal to decadal timescales. At the Spanish level, BSC stands as the national reference in this research field. The VOLCADEC project benefits from the strong experience of all the groups involved in SPECS, in particular KNMI (Netherlands), ECMWF and MetOffice (UK), Météo-France and CERFACS (France) and SMHI (Sweden). As stated in the previous summary, the experiments launched under the VOLCADEC project will serve as a base to involve the climate prediction team in DCPD and VOLMIP, two model intercomparison projects planned under CMIP6 that will reinforce our position at the international level.

Sets of experiments

The VOLCADEC project focuses on two major issues: (1) Verifying the ability of current climate models to forecast the climate response to volcanic eruptions and setting up new protocols to take into account stratospheric volcanic aerosols in climate forecast systems; (2) Using an original approach to investigate the regional climate impacts of volcanic aerosols. The first issue will be approached with a first set of experiments run with the version 2.3 of EC-Earth (Part 1), and the second one will be based on the analysis of a second set run with the version 3.2 of this model. Note that a one-year EC-Earth simulation need approximately 300 CPU hours on the Marenostrum3 BSC supercomputer.

#### Part 1: Forecasting the climate response to volcanic eruptions

Under the CMIP5 exercise (Taylor et al., 2005), the climate prediction team of the BSC took part in the DCCP project, by running 5-member decadal hindcasts, initialised in November of each year from 1960 to 2015. Here, we will run again these hindcasts for the periods covering the last large eruptions, but with a model configuration that excludes the volcanic forcing. This consists in simulations initialised the years surrounding the 1963 Agung, the 1982 el Chichón and the 1991 Pinatubo eruptions, i.e. fifteen 10-year and 5-member experiments, reaching a total of 750 years of simulation. This set of experiments will allow investigating the atmospheric and oceanic responses to the last eruptions in decadal hindcasts with the final goal to diagnose the forecast system skill that is related to the volcanic forcing. A second set of experiments will be conducted following a similar protocol, but with hindcasts using an idealized volcanic forcing that could be used for real-time forecast. This idealized forcing is based on an estimation of the observed volcanic forcing that decreases with an exponential decay during the forecast. It is representative as the protocol of a real-time forecast, since it does not use any information of the future (as retrospective hindcasts typically include observed forcings and cannot be considered as real forecasts). This second set consists in a total of 750 years of simulation. It will be completed with two 10-year and 20 member hindcasts initialised in 2014, one excluding and another one including a Pinatubo-like eruption (a total of 400 years). This last set is a common experiment required for the VolMIP and DCCP initiatives. It aims at comparing CMIP-model forecasts considering a potential new eruption that would have occurred recently. The first part of this RES application will allow producing experiments used to evaluate the ability of the EC-Earth forecast system to predict the climate response to volcanic eruptions. It is based on a total of 1900 years of simulation that requires a total of ~570000 hours CPU (~300\*1900).

#### Part 2: Investigating the regional climate impacts of a volcanic eruption.

Under the CMIP6 initiative, our team is currently contributing to the development of the last version of EC-Earth, by implementing in particular the forcing recommended under CMIP6. Our team is developing a new scheme for the stratospheric aerosols that includes a vertical resolution and a waveband dimension that was not included in the previous versions of this model. We expect this new parameterisation to allow a better representation of the particle interactions with the solar radiation. This new configuration will be used here to run a 1000-year control simulation including the pre-industrial forcing levels. Following the VolMIP protocol, we will then simulate a Pinatubo eruption initialised from climate conditions extracted from the control simulation and describing extreme states of the natural modes of

variability (ENSO, NAO, Atlantic and Pacific Multidecadal Variability, etc.). This experiment consists in a 10-year and 100-member simulation, reaching a total of 1000 years of simulation, which will be used to investigate how the climate conditions affect the climate response itself to a volcanic eruption. This approach has already been partly investigated by our team, questioning how the Atlantic Variability modulates the climate response to volcanic eruptions in the middle latitudes (Ménégoz et al., in rev). The second part of this RES project requires a total of 2000 years simulated with the version 3 of EC-Earth, reaching a cost of ~600000 hours (~300\*2000).

#### References:

- Bladé, I. et al., 2011, *Clim. Dyn.*, 39, 3-4.  
Doblas-Reyes, F., et al., 2013, *Wiley Interdisc. Rev.: Clim. Change*, 4(4), 245-268.  
Hansen, J. et al., 2011, *Atm. Chem and Phys*, 11(24), 13421-13449.  
Hawkins E., et al., 2009, *Bull. Amer. Meteor. Soc.*, 90(8), 1095-1107.  
Hazeleger, W. et al., 2010, *Bull. Amer. Meteor. Soc.*, 91, 1357–1363.  
Ménégoz, M. et al., in rev for *Clim. Dyn.*  
Solomon, S. et al., 2011, *Science*, 333(6044), 866–870.  
Swingedouw, [...], Ménégoz, M et al., *Glob. and Planet. Change*, Vol. 150, P. 24–45.  
Taylor, K. E. et al., 2012, *Bull. Amer. Meteor. Soc.*, 93, 485–498.

### **g) Computational algorithms and codes outline**

EC-Earth is a European model (<http://www.ec-earth.org/>) that comprises three major components: IFS (atmospheric component), NEMO (oceanic component) and OASIS3 (coupler). It is essential to configure and build separate executables for each one of them. IFS and NEMO fully support a parallel environment, while OASIS3 supports a pseudo-parallel environment. OASIS3 requires Cray pointers. For IFS there is a possibility to activate an OpenMP switch but in this case the implemented MPI should be thread-safe. IFS generates the output in GRIB format and NEMO in NetCDF, while OASIS3 does not generate any output. At the end of a simulation the three components always generate restarts separately (IFS in binary, and NEMO and OASIS3 in NetCDF format).

For configuring and building the model executable, GNU make 3.81 or 3.81+, FORTRAN 77/90/95 complaint compiler with preprocessing capabilities and NetCDF4 deployed with HDF5 and SZIP are needed. A newly designed tool for automatic build configuration called "ec-conf" can be used. This useful tool requires Python 2.4.3 or 2.4.3+ (although it does not work yet with Python 3.0+). For NEMO, FCM, bash and perl are essential, and the GRIB\_API I/O 1.9.9 or 1.9.9+ and GRIBEX 370 are required for IFS. GNU date (64-bit) is also required for executing the model with the run scripts.

Currently, several releases of EC-Earth are available, v2.3, and several versions of v3. This activity will be carried out with both versions v2.3 and v3.2. EC-Earth supports several configurations that have already been tested on various supercomputing platforms, Marenstrum3 among them. The experiments using the v3.2 in this activity will use the T255-ORCA1 configuration, which corresponds to a spatial resolution of 80 km in the atmosphere and 100 km in the ocean. In order to store sources and initial data, the experiments require at least ~100 GB of disk space for each release.

In the context of the experiments planned with EC-Earth v2.3, which will use the T159L62-ORCA1 configuration, a benchmarking performed previously suggests that, taking into account the average load of the Marenostrom queues, optimum performance is obtained when using 48 cores. This is a configuration which runs faster than previous experiments at Marenostrom but still remains scalable for this resolution. This configuration generates 2 GB of output per month of simulation.

### 3. Software and Numerical Libraries

---

Software components that the project team requires for the activity.

#### a) Applications + Libraries

BLAS, HDF5, LAPACK, SCALAPACK, UDUNITS

#### b) Compilers and Development Tools

GCC, INTEL, MVAICH2

#### c) Utilities + Parallel Debuggers and Performance Analysis Tools

CMAKE, PERL, PYTHON, VALGRIND, NCVIEW, AUTOCONF

#### d) Other requested software

GRIB\_API, GRIBEX, CDO, NCO and CDFTOOLS

#### e) Proprietary software

### 4. Research Team Description

---

#### a) Personal Data

<b>Name of Team Leader</b>	Martin Ménégos
<b>Institution</b>	Barcelona Supercomputing Center
<b>e-mail</b>	<a href="mailto:martin.menegoz@bsc.es">martin.menegoz@bsc.es</a>
<b>Phone</b>	+34 934137644
<b>Nationality</b>	France

#### b) The employment contract of the activity leader with the research organisation is valid at least 3 months after the end of the allocation period.

Yes

#### c) Curriculum Vitae of the Team Leader

Dr Martin Ménégos is the Principal Investigator (PI) of this proposal. He is a research

associate at the Earth Science department led by Prof. Doblas-Reyes of the BSC. Dr Ménégos received his PhD at Météo France (Toulouse) in 2009, where he studied the interactions between the atmosphere and natural and anthropogenic aerosols. After investigating the sinks and sources of dust, sulphate and black carbon, to evaluate the particle pollution rates at the global scale and more regionally in the Arctic and in Europe, he focused on their impacts on the climate system. He highlighted the links between aerosols and weather regimes in Europe.

Located in French CNRS institutes from 2010 to 2014, Dr Ménégos investigated the sensitivity of glaciers and snow cover to greenhouse gases and aerosol pollution, by using and developing both global and regional climate models. He modelled in particular how the aerosol deposition affects the snow cover during the last century and in different climate scenarios over the XXIth century. He has been involved in the evaluation of the CMIP5 models to simulate snow cover and its sensitivity to global warming. He also supervised field campaigns in various mountainous areas around the world to measure glaciers mass balance.

Now at BSC, Dr Ménégos leads the research line devoted to stratosphere-troposphere coupling processes in the Earth Sciences Department. He is largely involved in the development of the EC-Earth European model. He is applying this model in initialised simulations to make quasi-operational forecasts at seasonal-to-decadal timescales. He investigates the possibility to predict the climate response to large volcanic eruptions. He also takes part in a collaborative development of a statistical tool to assess the performance of forecast systems. He developed collaborations with 13 research institutes, organizing its research through several European and national projects, including one MINECO project that he is coordinating (VOLCADEC, 168Keuros). He has an H-index of 9 (Google Scholar), with 17 articles in peer-review journals (6 as first author), 3 articles submitted and 500 citations. He presented his results through 38 conferences and international project meetings. His scientific background and his skill in modelling coupled to his large international network of collaborations consist in solid basis to develop climate models and forecast systems. His full CV is available at <http://martinmenegoz.neowordpress.fr/>.

#### **d) Names of other researchers involved in this activity**

Virginie Guemas ([virginie.guemas@bsc.es](mailto:virginie.guemas@bsc.es))

Francisco Doblas-Reyes ([francisco.doblas-reyes@bsc.es](mailto:francisco.doblas-reyes@bsc.es))

Omar Bellprat ([omar.bellprat@bsc.es](mailto:omar.bellprat@bsc.es))

Eleftheria Exarchou ([eleftheria.exarchou@bsc.es](mailto:eleftheria.exarchou@bsc.es))

Chloé Prodhomme ([chloe.prodhomme@bsc.es](mailto:chloe.prodhomme@bsc.es))

Neven Fuckar ([neven.fuckar@bsc.es](mailto:neven.fuckar@bsc.es))

Francois Massonnet ([francois.massonnet@bsc.es](mailto:francois.massonnet@bsc.es))

Mario Acosta ([mario.acosta@bsc.es](mailto:mario.acosta@bsc.es))

Roberto Bilbao ([roberto.bilbao@bsc.es](mailto:roberto.bilbao@bsc.es))

Laurent Brodeau ([laurent.brodeau@bsc.es](mailto:laurent.brodeau@bsc.es))

Etienne Tourigny ([etienne.tourigny@bsc.es](mailto:etienne.tourigny@bsc.es))

Valentina Sicardi ([valentina.sicardi@bsc.es](mailto:valentina.sicardi@bsc.es))

**e) Relevant publications**

Ménégoz, M., Cassou, C., Swingedouw, D., Doblas-Reyes, F., Bretonnière, P.A., Modulation of the climate response to a Pinatubo eruption by the Atlantic Multi-decadal Variability, in revision for Climate Dynamics.

Siegert, S., Stephenson, D. B., Bellprat, O., Ménégoz, M., & Doblas-Reyes, F. J., 2016: Detecting improvements in forecast correlation skill: Statistical testing and power analysis. Monthly Weather Review, doi: <http://dx.doi.org/10.1175/MWR-D-16-0037.1>

Guemas V., F.J. Doblas-Reyes, I. Andreu-Burillo and M. Asif (2013). Retrospective prediction of the global warming slowdown in the past decade. Natuer Climate Change, 3, 649-653, doi:10.1038/nclimate1863.

Doblas-Reyes, F. J., Andreu-Burillo, I., Chikamoto, Y., García-Serrano, J., Guemas, V., Kimoto, M., ... & Van Oldenborgh, G. J. (2013). Initialized near-term regional climate change prediction. Nature communications, 4, 1715.

Ménégoz, M., Guemas, V., Salas y Melia, D. and Voltaire, A, 2010: Winter interactions between aerosols and weather regimes in the North-Atlantic European region, J. Geophys. Res., 115, D09201, doi:10.1029/2009JD012480, <http://onlinelibrary.wiley.com/doi/10.1029/2009JD012480/abstract>

5. Resources

---

**a) Estimated resources required for the Activity for the current Application Period**

<b>Requested machine</b>	Finis Terrae II ((BULL with 7712 cores Intel Haswell 2680v3, Infiniband Fat-Tree FDR, 44TB memory, 1 PB disk)
<b>Interprocess communication</b>	Null

**Typical Job Run**

<b>Number of processors needed for each job</b>	256.00
<b>Estimated number of jobs to submit</b>	375.00
<b>Average job durations (hours) per job</b>	6.00
<b>Total memory used by the job (GBytes)</b>	100.00

**Largest Job Run**

<b>Number of processors needed for each job</b>	
<b>Estimated number of jobs to submit</b>	



**Average job durations (hours) per job****Total memory used by the job (GBytes)**

<b>Total disk space (Gigabytes)</b>	<b>Minimum</b>	8000.00	<b>Desirable</b>	10000.00
-------------------------------------	----------------	---------	------------------	----------

<b>Total scratch space (Gigabytes)</b>	<b>Minimum</b>	15000.00	<b>Desirable</b>	15000.00
--	----------------	----------	------------------	----------

<b>Total tape space (Gigabytes)</b>	<b>Minimum</b>	0.00	<b>Desirable</b>	0.00
-------------------------------------	----------------	------	------------------	------

<b>Total Requested time (Thousands of hours)</b>		576.00		
--	--	--------	--	--

If this activity is asking for more than 2Million CPU hours, you need to justify the amount of resources requested for the activity. (max 1000 characters)

**INFORMATION:** The estimated cost of the requested hours, considering only the electricity cost, is 6583.68 euros.

The architectures selected for the requested resources are only a suggestion. If no hours in this machine/these machines are available, please grant resources in any other similar architecture where the codes used for the application may run efficiently.

\*\* this option implies that if no hours in this machine/these machines are available, the acces committee will reject the full application.

**b) Estimate of the total resources that the Activity will require until it is completed (including the present and all the following Application Periods)**

**Number of application periods expected to complete this Activity**

1

**Total Requested Time (thousands of hours) expected to complete this Activity (sum of both periods)**

1176.00

## 6. Abstract for publication

---

Martin Ménégoz is coordinating at BSC the "Volcanic activity in seasonal to decadal climate forecasts" project, (VOLCADEC) with the aim to answer how forecast systems should take into account stratospheric particles. It focuses on two main challenges: (1) how can the climate response to a large volcanic eruption and to stratospheric aerosols changes be predicted? (2) How climate conditions modulate the climate response to large volcanic eruptions? The final goal of VOLCADEC is to design the optimal strategy to take into account stratospheric aerosols in climate forecast systems. Optimized systems will be very helpful to

provide climate information, in particular in the Mediterranean region, where the climate variability is largely dependent on the stratospheric aerosols load.

## 7. Contact with CURES during last year

---

Information about the RES Users Committee (CURES).

### **a) User has contacted the CURES during last year**

No

### **b) If not, indicate why you have not contacted the CURES**

Because this is my first application to RES.

Barcelona Supercomputing Center, 2016